

EXERCISE 1

Design ϵ -NFA's for the following languages.

- The set of strings consisting of zero or more a's followed by zero or more b's, followed by zero or more c's.
- The set of all strings that consist of either 01 repeated one or more times or 010 repeated one or more times.

EXERCISE 2



Consider the following ϵ -NFA's.

	ϵ	a	b	c	
$\rightarrow P$	$\{q, r\}$	$\emptyset$	$\{q\}$	$\{r\}$	} ϵ -NFA ₁
q	$\emptyset$	$\{P\}$	$\{r\}$	$\{P, q\}$	
* r	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	
$\rightarrow P$	$\emptyset$	$\{p\}$	$\{q\}$	$\{r\}$	} ϵ -NFA ₂
q	$\{p\}$	$\{q\}$	$\{r\}$	$\emptyset$	
* r	$\{q\}$	$\{r\}$	$\emptyset$	$\{p\}$	


- Compute the ϵ -closure of each state.
- Give all the strings of length three or less accepted by the automata.
- Convert the automata to DFA's.

EXERCISE 3


Convert the following ϵ -NFA's to a DFA's.



1) a)



1) b)



2) a) Remember that: $\begin{cases} q \in \text{ECLOSE}(q) \\ p \in \text{ECLOSE}(q) \ \& \ r \in \delta(p, \epsilon) \Rightarrow r \in \text{ECLOSE}(q) \end{cases}$

$\epsilon\text{-NFA}_1$: $\text{ECLOSE}(p) = \{p, q, r\}$, $\text{ECLOSE}(q) = \{q\}$, $\text{ECLOSE}(r) = \{r\}$

$\epsilon\text{-NFA}_2$: $\text{ECLOSE}(p) = \{p\}$, $\text{ECLOSE}(q) = \{p, q\}$, $\text{ECLOSE}(r) = \{p, q, r\}$

2) b) $\epsilon\text{-NFA}_1$

All strings except for: bba, bbb, bbc

$\epsilon\text{-NFA}_2$

All strings except for: $\epsilon, a, b, aa, ab, ba, aaa, aab, aba, baa$

2) c) From $\epsilon\text{-NFA}$ to NFA

Remember that $\delta_N(q, a) = \hat{\delta}_\epsilon(q, a) = \text{ECLOSE}\left(\bigcup_{p \in \text{ECLOSE}(q)} \delta(p, a)\right)$

2) c) can't

E5.3

The calculations for ε -NFA₁ go as follows:

$$\hat{\delta}_\varepsilon(p, a) = \text{ECLOSE}(\delta(p, a) \cup \delta(q, a) \cup \delta(r, a)) = \text{ECLOSE}(\emptyset \cup \{p\} \cup \emptyset) = \{p, q, r\}$$

$$\hat{\delta}_\varepsilon(p, b) = \text{ECLOSE}(\delta(p, b) \cup \delta(q, b) \cup \delta(r, b)) = \text{ECLOSE}(\{q\} \cup \{r\} \cup \emptyset) = \{q, r\}$$

$$\hat{\delta}_\varepsilon(p, c) = \text{ECLOSE}(\delta(p, c) \cup \delta(q, c) \cup \delta(r, c)) = \text{ECLOSE}(\{r\} \cup \{p, q\} \cup \emptyset) = \{p, q, r\}$$

$$\hat{\delta}_\varepsilon(q, a) = \text{ECLOSE}(\delta(q, a)) = \text{ECLOSE}(\{p\}) = \{p, q, r\}$$

$$\hat{\delta}_\varepsilon(q, b) = \text{ECLOSE}(\delta(q, b)) = \text{ECLOSE}(\{r\}) = \{r\}$$

$$\hat{\delta}_\varepsilon(q, c) = \text{ECLOSE}(\delta(q, c)) = \text{ECLOSE}(\{p, q\}) = \{p, q, r\}$$

$$\hat{\delta}_\varepsilon(r, a) = \text{ECLOSE}(\delta(r, a)) = \text{ECLOSE}(\emptyset) = \emptyset$$

$$\hat{\delta}_\varepsilon(r, b) = \text{ECLOSE}(\delta(r, b)) = \text{ECLOSE}(\emptyset) = \emptyset$$

$$\hat{\delta}_\varepsilon(r, c) = \text{ECLOSE}(\delta(r, c)) = \text{ECLOSE}(\emptyset) = \emptyset$$

By doing the same calculations for ε -NFA₂ we get the following NFA's

δ_N	a	b	c	
$\xrightarrow{*} p$	$\{p, q, r\}$	$\{q, r\}$	$\{p, q, r\}$	NFA ₁ (note that ε -NFA ₁ accepts the empty string ε)
q	$\{p, q, r\}$	$\{r\}$	$\{p, q, r\}$	
$* r$	$\emptyset$	$\emptyset$	$\emptyset$	
$\rightarrow p$	$\{p\}$	$\{p, q\}$	$\{p, q, r\}$	NFA ₂
q	$\{p, q\}$	$\{p, q, r\}$	$\{p, q, r\}$	
$* r$	$\{p, q, r\}$	$\{p, q, r\}$	$\{p, q, r\}$	

From NFA to DFA

This was already done in exercise E4 (06/11/2008)

3) ϵ -NFA₁

E5.4

From ϵ -NFA₂ to NFA₁ (as above)

δ_{NFA_1}	x	y	z
$\xrightarrow{*} p$	$\{p, q, r\}$	$\{q, r\}$	$\{r\}$
q	$\emptyset$	$\{q, r\}$	$\{r\}$
$* r$	$\emptyset$	$\emptyset$	$\{r\}$

From NFA₁ to DFA₁ (subset construction)

δ_D	x	y	z
$A = \emptyset$	A	A	A
$\xrightarrow{*} B = \{p\}$	H	G	D
$C = \{q\}$	A	G	D
$* D = \{r\}$	A	A	D
$* E = \{p, q\}$	H	G	D
$* F = \{p, r\}$	H	G	D
$* G = \{q, r\}$	A	G	D
$* H = \{p, q, r\}$	H	G	D

ϵ -NFA₂

We only provide the final DFA

